

Bert Cregg and Robert Schutzki
Michigan State University
Department of Horticulture and Department of Forestry





Emerald Ash Borer

his guide gives suggestions for species that should be considered in situations where a homeowner, landscaper, or urban forester may have planted an ash in the past. General species characteristics, mature form and size, and other noteworthy characteristics are given; all are hardy to at least zone 5. Diversity of species is also important. Look around you, and plant a variety trees.

'Autumn blaze' Freeman maple Acer x freemanii

Height: 50-55 ft. **Spread:** 30-40 ft.

Site preferences: Zone 4-7. Prefers moist soil but is drought tolerant. pH: adaptable. Fast growing.


Ornamental characteristics: Shape:

oval-rounded. Foliage has excellent orange-red fall color.

Insect/disease factors: None

serious.

Cultivars: Marmo, Sienna Glen



Trident maple Acer buergeranum

Height: 20-25 ft. **Spread:** 20-35 ft.

Site preferences: Zone 5b or 6a-8a. Prefers full sun.
Adapted to urban conditions. Moderate salt tolerance. Tolerates moderate drought. pH: 5.0-7.0.



Ornamental characteristics:

Shape: round. Mottled bark, shedding with age; dense, dark green leaves turning orange to red in the fall. Introduced from China.

Insect/disease factors: None serious.

Hedge maple Acer campestre

Height: 25-30 ft. **Spread:** 25-30 ft.

Site preferences: Zone 5a-8b. Sun to partial shade. Tolerates drought. Moderate salt tolerance. pH: 5.0-8.2. Adapted to compaction, severe pruning.



Ornamental

characteristics: Shape: round. Excellent residential use under utility lines. Easily transplanted in spring or fall. Cultivars: 'Queen Elizabeth' (more upright, vigorous [zone 6b-7b]); 'Schichtel's Upright' (5b-8b). Introduced from Europe, Near East and Africa.

Insect/disease factors: Occasional problems with

verticillium wilt.

Miyabe maple Acer miyabei

Height: 30-40 ft. **Spread:** 20-30 ft.

Site preferences: Zone 4-7. Moist, well-drained soil. pH:

adaptable.


Ornamental characteristics:

Shape: rounded.

Insect/disease factors: None

serious.

Cultivars: 'State Street' —
fast growing, upright oval
habit, dark green leaves
with yellow fall color, corky bark.






Norway maple Acer platanoides

Height: 40-50 ft. **Spread:** 30-50 ft.

Site preferences: Zone 4-7. Prefers moist soils and will tolerate clay (but not wet) soil. pH: very adaptable.

Ornamental characteristics:

Shape: rounded. Foliage is dark green with occasionally good yellow fall color.



Insect/disease factors:

Verticillium wilt and anthracnose.

Cultivars: 'Crimson King' — reddish purple foliage; 'Columnare' — narrow oval habit (20-ft. spread); 'Summershade' — fast growing, heat resistant; 'Easy Street' — narrow oval.


Red maple Acer rubrum

Height: 40-60 ft.

Site preferences: Zone 3b-9a. Moist soils necessary; flood tolerance of specific cultivars is listed below. Salt sensitive. pH: 5.0-7.0 (chlorosis occurs at high pH).


Ornamental characteristics:

Shape: round, upright. Good fall color.



Leafhopper and borers.

Cultivars: 'Autumn Flame' (early, low flood tolerance, persistent red fall color, zone 3b); 'Northwood' (low flood tolerance, good orange-red fall color, zone 3b); 'October Glory' (intermediate flood tolerance, excellent red fall color, zone 5a); 'Red Skin' (large, thick foliage, early reddish maroon fall color, rare, zone 4); 'Red Sunset'. Transplant in spring, high flood tolerance, specify "own-rooted" — graft incompatibility can be a problem. 'Redpointe' — pyramidal with dominant central leader.




Sugar maple Acer saccharum

Height: 60-80 ft. **Spread:** 40-50 ft.

Site preferences: Zone 4-8. Prefers moist, well-drained soil. pH: neutral to slightly

Ornamental characteristics:

Shape: oval to rounded. Good fall color of yellow, orange or red.




Insects/disease factors: verticillium wilt, leaf scorch. Cultivars: 'Crescendo' — dark green foliage with good orange-red fall color, heat and drought tolerant; 'Green Mountain' — thick, dark green leaves that resist leaf scorch, yellow fall color; 'Legacy' — thick and glossy dark green leaves, drought tolerant, yellow-orange fall color.

Shantung maple Acer truncatum

Height: 25-30 ft. **Spread:** 20-30 ft.


Site preferences: Zone 5a-7b (species is variable; hybrid cultivars, zone 5). Requires moist, well-drained site; tolerates moderate drought. Moderate salt tolerance. pH: 5.0-8.2. Resistant to leaf scorch.



Ornamental characteristics:

Shape: round. Dark green, glossy leaves with early fall color of yellow-orange to bright red.

Cultivars: Hybrids of A. platanoides are 'Norwegian Sunset' (upright round, orange-red to red fall color, more heat and drought tolerant than A. platanoides) and 'Pacific Sunset' (upright oval, finer branched



Insect/disease factors: None serious.



Horse chestnut Aesculus hippocasteana

Height: 50 ft. -75 ft. Spread: 40 ft. -70 ft. Site preferences: Zone 4-8. Prefers moist, welldrained soils. pH: adaptable.

Ornamental characteris-


tics: Shape: rounded. Other: showy white flowers in mid-May.

Insect/disease factors:

Leaf scorch can occur under hot/dry conditions in late summer.

Cultivars: 'Baumanii' more showy with longer lasting

flowers and seedless.



Upright European hornbeam Carpinus betulus 'Fastigiata'

Height: 35 ft. Spread: 25 ft.


Site preferences: Zone 5a-7a. Tolerates drought, heavy soil. Salt sensitive. pH: 5.0-8.2.

Ornamental characteristics:

Shape: narrow when young, becoming oval; narrow branching angles.

Insect/disease factors: Canker, leaf spot and twig blight can occur if the tree is stressed.

Additional: Transplant in spring; somewhat slow to establish from bare root.



American hornbeam Carpinus caroliniana

Height: 20-30 ft. Spread: 20-30 ft. Site preferences: Zone 3b-9a. Prefers moist soil but will tolerate some intermittent drought. pH: 5.0-8.2. Prefers shaded, moist soils.

Ornamental characteris-

tics: Shape: round,

spreading. "Muscled" bark, good orange-red fall color, clumps or single stem.

Insect/disease factors: Relatively pest and disease free, requires little maintenance, though damage by ice storms can be a problem. Cankers can occur with specimens of southern origin, so it is important to know the source of the tree.

Additional: Transplant in spring; somewhat slow to establish. Native to Michigan and eastern United States.



Height: 40-60 ft. Spread: 30 ft.

Site preferences: Zone 3b-8b. Tolerates drought. Salt sensitive. pH: 5.0-8.2. Tolerates light shade, wind, heat.

Ornamental characteris-

tics: Shape: pyramidal when young; open, irregular when mature, with dark green, leathery leaves.

Insect/disease factors:


Witch's broom and insect

galls (aesthetic but not life-threatening problems).

Cultivars: 'Prairie Pride' (rapid, compact grower; thick, leathery foliage).

Additional: Transplant in spring; somewhat slow to

establish.





Katsura tree Cercidiphyllum japonicum

Height: 40 ft. Spread: 40 ft.

Site preferences: Zone 5a-9a. Requires evenly moist soils; not drought tolerant. pH: 5.0-8.2. Does not tolerate compaction or heavy soils; tolerates light shade. Protected site best.


Ornamental characteris-

tics:

Shape: upright, pyramidal when young, becoming rounded with age.

New leaves are bright red when unfolding; good yellow fall color; dry leaves have caramel scent; clump or single

Insect/disease factors: None serious.



Yellowwood Cladrastis kentukea

Height: 30 ft. Spread: 40 ft. Site preferences: Zone 4b-8a. Best in

moist, well- drained soils; not drought tolerant. pH: 5.0-8.2.

Ornamental characteristics: Shape: round. Bright green foliage; showy, fragrant flowers in

late spring. Insect/disease factors: None.

Additional: Prune only in summer — profuse bleeding will occur at other times of the year. Specify single leader can be weak-wooded with poor crotch angles that split out as the trees grow older. Transplant in spring.

Turkish filbert Corylus colurna


Height: 40-50 ft. Spread: 20-30 ft. Site preferences: Zone 4-7. Drought tolerant once established. pH:

adaptable.

Ornamental character-


istics: Shape: broad pyramidal, maintains formal shape. Other: attractive dark green

foliage.



None serious.

Additional: This is a low-maintenance but attractive species that is quite underused.



Hardy rubber tree Eucommia ulmoides

Height: 40-60 ft. Spread: 45 ft.

Site preferences: Zone 5b-8.2. Tolerates drought; intolerant of poor drainage. pH: 5.0-8.2. Tolerates heat; needs full sun.


Ornamental characteris-

tics: Shape: round, wide, spreading with ascending branches. Glossy green foliage.

Insect/disease factors:

None.

Additional: Transplants easily in spring or fall.





Gingko Gingko biloba

Height: 40-50 ft.

Spread: Variable, depending

on cultivar.

Site preferences: Zone 4b-8b. Tolerates drought. Moderate salt tolerance.

pH: 5.0-8.2.

Ornamental characteristics:

Shape: irregular when young, becoming pyramidal with age. Yellow fall color.

Cultivars: 'Autumn Gold' (symmetrical, broad, out-



Additional: Specify male trees because of noxious fruit smell (all cultivars are male). Transplant in spring or fall.

Insect/disease factors: None serious.



Height: 40-100 ft. **Spread:** 40 ft.

Site preferences: Zone 4b-9a. Tolerates wet and dry sites. High salt tolerance.

pH: 5.0-8.2.

Ornamental characteristics:

Shape: Open, spreading.


Insect/disease factors:

Overplanting has encouraged severe insect problems in many areas (honeylocust

plant bug, spider mite, borer, webworm).

Cultivars: 'Shademaster' (high, vase-shaped canopy, essentially fruitless); 'Skyline' (upright, pyramidal, strong central leader, good fall color); 'Northern Acclaim' (most cold hardy, good form).

Additional: Transplants easily in spring. 'Inermis' refers to the thornless selection.



Kentucky coffee tree Gymnocladus dioicus

Height: 60-75 ft.

Spread: 40-50 ft.


Site preferences: Zone

3b-8. Drought tolerant after establishment.

pH: adaptable.

Ornamental characteris-

tics: Shape: oval crown with irregular branching. Foliage is dark green with compound leaves; can sometimes have a good yellow fall color.



Insect/disease factors: None serious.

Cultivars: 'Prairie Titan' — nice branching habit and bluegreen summer foliage. 'Espresso' — seedless, good form.

Sweetgum Liquidambar styraciflua

Height: 70-100 ft. **Spread:** 40-65 ft.

Site preferences: Zone 5b-9a. Tolerates intermittent flooding, moderate drought. pH: 5.0-7.5. Best on moist,


sunny sites.

Ornamental characteris-

tics: Shape: pyramidal when young, becoming round with age.

Excellent fall color. Fruit litter may be

objectionable.



Insect/disease factors: None serious.

Cultivars: 'Moraine' (most cold hardy, zone 4b, bright red fall color); 'Rotundaloba' (fruitless, not as hardy); 'Worplesdon' (orange-red fall color, broadly pyramidal to oval).

Additional: Northern seed source recommended. Transplant in spring. Iron chlorosis can occur.



Tulip tree Liriodendron tulipifera

Height: 80-120 ft. Spread: 35-50 ft.

Site preferences: Zone 5a-9a. Requires evenly moist soils but does not tolerate poor drainage. Needs adequate space. Salt sensitive.

pH: 5.0-8.2.


Ornamental characteristics:

Shape: broad pyramidal. Yellow fall color. Somewhat weak-wooded.

Insect/disease factors:

Occasional problems with leaf scorch, canker, and verticillium wilt.

Additional: Transplant in spring. Northern seed source recommended.



Dawn redwood Metasequoia glyptostroboides

Height: 70-100 ft. Spread: 25 ft.

Site preferences: Zone 5b-9a. Requires moist sites, will not tolerate dry sites. Salt sensi-

tive. pH: 5.0-7.5.

Ornamental characteristics:

Shape: pyramidal. Lower branches need to be removed for streetside use.


Fast growing.

Insect/disease factors:

Relatively trouble free, though Japanese beetle can

be a nuisance.

Additional: Transplant in spring or fall.



Amur maackia Maackia amurensis


Height: 20-30 ft. Spread: 30 ft. Site preferences: Zone 4-7. Moist, well-drained soil.

pH: adaptable. Ornamental charac-

teristics: Shape: rounded. White flowers in June; copper-brown, peeling, attractive

bark.

Insect/disease factors: None serious.



Tupelo Nyssa sylvatica

Height: 35 ft. Spread: 20 ft.

Site preferences: Zone 5a-9a. Tolerates wet sites, intermittent drought. Relatively salt tolerant.

pH: 5.5-7.0.

Ornamental characteris-

tics: Shape: pyramidal. Bright red fall color.


Insect/disease factors:

None.

Additional: Difficult to transplant — use small sizes, balled and

burlapped only. Slow to recover from transplanting.

Transplant in spring.





American hophornbeam Ostrya virginiana

Height: 40 ft. Spread: 25 ft.

Site preferences: Zone 3b-9a. Prefers moist soil but will tolerate moderately dry soils once established. Very salt sensitive. pH: 5.0-8.2. Tolerates light shade.



Shape: upright oval. Foliage is dark green; fall color is yellow. Slow growing.

Insect/disease factors: None.

Additional: Slow to recover from transplanting. Transplant

in spring.




Height: 30-45 ft. Spread: 40-50 ft. Site preferences: Zone 3b-7. Drought and pollution tolerant. pH: adaptable.


Ornamental characteristics: Shape: broad, rounded. Yellow to bronze-yellow fall color. Ridged, corky bark in

old age.

Insect/disease factors: None serious.

Cultivars: 'Eye Stopper' 'His Majesty' 'Macho'





London planetree Platanus x acerifolia

Height: 50 ft. Spread: 40 ft. Site preferences:


Zone 5b-9a. Tolerates wet and intermittently

flooded sites;

drought tolerant. Salt tolerant. pH: 5.0-8.2.

Ornamental charac-

teristics: Shape: pyramidal when



young, becoming open, spreading with age. Medium green foliage, exfoliating bark.

Insect/disease factors: Overplanting has encouraged disease and insect problems in New York City. 'Liberty' resistant to anthracnose and powdery mildew; 'Bloodgood' moderately resistant. Cankerstain may be a problem.

Cultivars: 'Bloodgood' (maple-green foliage; tolerates

severe pruning; rapid grower); 'Liberty'. Additional: Transplant in spring or fall.

Common chokecherry Prunus virginiana

Height: 20-30 ft. **Spread:** 18-25 ft.

Site preferences: Zone 2-6. Tolerant of moist and dry

sites.

pH: adaptable.


Ornamental

characteristics: Shape: rounded. White flowers in early May, red fruit that ripen to a dark purple.

Insect/disease factors:

Black knot.

Cultivars: 'Schubert' and 'Canada Red' — foliage emerges green but changes to a reddish purple for the remainder of the growing season; pyramidal habit.





Callery pear Pyrus calleryana

Height: 50 ft. Spread: 30 ft. Site preferences:

Zone 5-8. Tolerant of moist and dry sites. pH: adaptable.

Ornamental


characteristics: Shape: formal rounded pyramidal. Foliage is glossy dark green; good red to purplish fall color. White flowers in late April or



Insect/disease factors: Fire blight.

Cultivars: 'Aristocrat' — wider shape, yellow to red fall color; 'Red spire' — pyramidal habit, yellow fall color;

'Chanticleer' — oval to narrow oval.



Swamp white oak Quercus bicolor


Height: 45 ft. Spread: 45 ft.

Site preferences: Zone 4a-8b. Tolerates temporary flooding, wet soils and somewhat dry soils. Salt sensitive. pH: 5.0-


7.0.

Ornamental

characteristics: Shape: broad oval with round top. Foliage is green with wavy margins.



None serious. Variable susceptibility to iron chlorosis.





Sawtooth oak Quercus acutissima


Height: 40 ft. Spread: 40 ft.

Site preferences: Zone 5b or 6a-9a. Tolerates moderately dry soils. Moderate salt tolerance. pH: 5.0-7.5. Good growth rate in moist sites.

Ornamental characteris-

tics: Shape: rounded. Foliage is dark green, glossy. Fall color is yellow.

Insect/disease factors: Reasonably pest free; preferred by orange-striped oakworm.





Northern pin oak Quercus ellipsoidalis

Height: 50-60 ft. Spread: 50 ft. Site preferences:

Zone 4-6. Prefers moist, well-drained soil but tolerates wet and dry conditions. pH: neutral to slightly acidic.

Ornamental

characteristics: Shape: pyramidal. Can have nice fall color.

Insect/disease factors:

Stem and leaf galls.

Additional: Does not have the chlorosis problems that occur in pin oak.






Shingle oak Quercus imbricaria

Height: 50 ft. Spread: 40 ft. Site preferences:

Zone 5a-9a. Prefers moist soil but will tolerate intermittent drought. pH: 5.0-7.5.

Ornamental

characteristics: Shape: broadly oval. Foliage is dark green, glossy. Fall color is yellowish to rusty red.



Insect/disease factors: Anthracnose, canker, powdery mildew, rust, wilt, galls, scales, borers, miners, etc., may be a problem.

Additional: Slow to recover from transplanting. Transplants well in spring.

Bur oak Quercus macrocarpa

Height: 55 ft. Spread: 45 ft.

Site preferences: Zone 3a-9a. Tolerates drought and intermittent flooding.

pH: 5.0-8.2.

Ornamental characteristics:


Shape: broadly oval, irregular and open. Foliage is dark green. Fall color is yellow to yellow-brown. Corky bark.



Insect/disease factors:

Susceptible to anthracnose, intermediate preference by orange-striped oakworm but generally not seriously affected.

Additional: Slow growing, massive tree needs adequate space. Transplant small sizes in spring.





Chinkapin oak Quercus muehlenbergii


Height: 45 ft. Spread: 45 ft.

Site preferences: Zone 5a-8b. Tolerates moderate drought. pH: 5.0-8.2. Extremely tolerant of alkaline soil.

Ornamental

characteristics: Shape: rounded, open. Foliage is lustrous dark green. Fall color is yellow to yel-

low-brown.



Insect/disease factors: None serious. Additional: Transplant in spring.

Japanese pagodatree Sophora japonica


Height: 50-70 ft. **Spread:** 40-70 ft. Site preferences:


Zone 5b-9a. Tolerates drought. Salt tolerant.

pH: 5.0-8.2.

Ornamental character-

istics: Shape: dense, round. Attractive cream-colored flowers in late July.





'Regent' resistant to leafhoppers, twig dieback and stem canker in colder zones.

Cultivars: 'Princeton Upright' (upright); 'Regent' (rapid growth, profuse flowering, resistant to leaf chewers).

Additional: Fixes own nitrogen. Transplant in spring or fall.

Somewhat weak wooded.



Linden or Basswood Tilia americana


Height: 35-50 ft.

Spread: 25-30 ft.

Site preferences: Zone
3a-8a. Requires moist,
well-drained sites but
will tolerate intermittent
drought once
established. Salt sensitive. pH: 6.5-8.2.

Ornamental characteris-

tics: Shape: narrow to broad pyramidal, depending on cultivar.



Foliage is light green to dark green, depending on cultivar. Fragrant flowers in June.

Insect/disease factors: Very susceptible to Japanese beetles; mites, aphids, borers, leaf miners and scale may also be problematic.

Cultivars: 'Boulevard' (narrowly pyramidal, hardy to zone 3); Legend® (broadly pyramidal, zone 4, medium green); 'Redmond' (pyramidal, light green foliage).

Additional: Transplant in spring or fall.

Little-leaf linden Tilia cordata

Height: 60-70 ft.

Spread: 40-50 ft.

Site preferences:

Zone 3b-7. Prefers


moist, well-drained soil.

pH: adaptable.

Pollution tolerant.

Ornamental characteris-

tics: Shape: rounded and densely branched. Yellowish, fragrant flowers that emerge in late June.



Insects/disease factors: Japanese beetle, aphids, scab and linden mite.

Cultivars: 'Greenspire' — single leader and nice branching form; 'June Bride' — more formal habit and greater flowering; 'Harvest Gold' — very hardy, bright yellow fall color.

Silver linden Tilia tomentosa

Height: 50-70 ft. Spread: 30-50 ft. Site preferences:

Zone 3b-7. Prefers moist, well-drained soils but is somewhat drought and heat tolerant.

pH: adaptable.

Ornamental character-

istics: Shape: upright oval. Glassy dark green leaves that are silvery underneath.

Yellowish white, fragrant flowers in early July.

Insect/disease factors: Same as little-leaf linden but less

susceptible.

Cultivars: 'Green Mountain' — fast growing, better drought and heat tolerance; 'Sterling' — broad pyramidal habit, resistant to Japanese beetle.



Elm hybrids Ulmus spp.

Height: 40-50 ft. **Spread:** 30-50 ft.

Site preferences: Zone varies. Tolerates intermittent flooding and drought once established. pH: 5.0-8.2.

Ornamental characteristics: Shape: upright

vase.

Cultivars: Cultivars reportedly resistant to Dutch elm disease:

Hybrids: 'Accolade'

'Commendation'

'Frontier'

'Homestead'


'New Horizon'

'Triumph'

American: 'Valley Forge'

'Princeton'

'Jefferson'



Japanese zelkova Zelkova serrata

Height: 50-80 ft. Spread: 40-50 ft. Site preferences:

Zone 5. Prefers moist, well-drained soil but is drought tolerant. pH:

adaptable.


Ornamental characteristics: Shape: vaseshaped. Foliage has rusty orange fall color. Flaky, attractive bark.

Insect/disease factors:

None serious.


Cultivars: 'Green Vase' — fast growing, nice branching

habit, good fall color.



Thanks also to Keith Warren of J. Frank Schmidt & Son Company for his expertise in developing this publication.







MSU is an affirmative-action, equal-opportunity institution. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital status, or family status. • Issued in furtherance of Extension work in agriculture and home economics, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Thomas G. Coon, Extension director, Michigan State University, E. Lansing, MI 48824. • This information is for educational purposes only. References to commercial products or trade names do not imply endorsement by MSU Extension or bias against those not mentioned. This bulletin becomes public property upon publication and may be printed verbatim with credit to MSU. Reprinting cannot be used to endorse or advertise a commercial product or company.